

The $^{13}\text{C}/^{12}\text{C}$ of Atmospheric Methane

PAUL QUAY, JOHN STUTSMAN, AND DAVID WILBUR
School of Oceanography, University of Washington, Seattle, 98195

INTRODUCTION

For the past 6 years we have been measuring the $^{13}\text{C}/^{12}\text{C}$ of atmospheric CH_4 on air samples collected at three CMDL sites (BRW, MLO, and SMO) and on the Washington coast at 48°N 126°W .

The $^{13}\text{C}/^{12}\text{C}$ of atmospheric CH_4 is a tracer that can distinguish between CH_4 input from bacterial and non-bacterial CH_4 sources. Bacterial CH_4 is microbially produced in anoxic environments like swamps, bogs, rice paddies, and the rumens of cows. Non-bacterial CH_4 sources include thermogenically produced natural gas and CH_4 produced during the incomplete oxidation of plant material during biomass burning. Bacterial CH_4 has a $\delta^{13}\text{C}$ of about -60‰ (versus PDB) whereas the $\delta^{13}\text{C}$ of natural gas and CH_4 produced from biomass burning are about -40 and -27‰ , respectively [Quay *et al.*, 1991].

The spatial and temporal variations in the $^{13}\text{C}/^{12}\text{C}$ of atmospheric CH_4 depend on the variations of the relative strength and $^{13}\text{C}/^{12}\text{C}$ of the CH_4 sources and sinks. Over interannual time scales the trends in the $^{13}\text{C}/^{12}\text{C}$ of atmospheric CH_4 indicate changes in the source composition, i.e., the relative strength of bacterial versus non-bacterial CH_4 sources. Because CH_4 will likely contribute about 15% of the radiative forcing during the next century [Wigley and Raper, 1992], it is important to quantify the strength of the individual CH_4 sources, currently known to about $\pm 50\%$ and to determine whether the CH_4 source strengths are changing with time. This latter point has been underscored by the recently observed slowdown in the rate of CH_4 increase in the atmosphere [Dlugokencky *et al.*, 1994].

METHODS

The air samples are collected at approximately 2-week intervals, using pre-evacuated stainless steel flasks either 15- or 30-L in volume. The CH_4 is extracted from air in our laboratory using the procedure developed by Stevens and Rust [1982]. Briefly, the air is metered into a high-vacuum extraction line through a series of liquid nitrogen traps to remove H_2O , CO_2 , and N_2O . The air then passes through a bed of Schutze's reagent, I_2O_5 on silica, to oxidize CO to CO_2 that is trapped cryogenically. Then the CH_4 in the air is combusted over platinized silica at 800°C to CO_2 which is then trapped cryogenically. The yield of the procedure, determined from standards, is $100 \pm 2\%$ ($n = 114$). The $^{13}\text{C}/^{12}\text{C}$ of the CO_2 derived from CH_4 is measured on a Finnigan MAT 251 gas ratio isotope mass

spectrometer. The overall measurement precision is about $\pm 0.1\text{‰}$. We obtain a $\delta^{13}\text{C}$ of -41.73‰ (versus PDB) for NBS-16.

RESULTS AND DISCUSSION

The seasonal cycle in the $\delta^{13}\text{C}$ of CH_4 is greatest at 71°N (BRW), with an amplitude of $\sim 0.6\text{‰}$, and decreases southward to 14°S (SMO) where we measure no significant seasonal trend, i.e. $< 0.1\text{‰}$ (Figure 1). The seasonal trends at 71°N and 48°N can be approximated roughly by a single harmonic with an annual period. Episodes of high CH_4 concentrations associated with very depleted $\delta^{13}\text{C}$ values occur at these two sites in September-October of each year and are due to input of bacterial methane. Generally, at the northern hemisphere sites the lowest $\delta^{13}\text{C}$ values occur in the fall and the highest values occur in the summer. This trend toward higher summertime $\delta^{13}\text{C}$ values is expected if CH_4 oxidation by OH primarily controls the seasonal cycle because the $^{12}\text{CH}_4$ molecules react at a slightly faster rate ($1.0054\times$) than the $^{13}\text{CH}_4$ molecules [Cantrell *et al.*, 1990]. The annual mean $\delta^{13}\text{C}$ values increase southward from about -47.8‰ at 71°N to -47.3‰ at 14°S .

We calculate a global average $\delta^{13}\text{C}$ of CH_4 of approximately -47.5‰ . The mean global $\delta^{13}\text{C}$ value, when combined with the ^{14}C content of atmospheric CH_4 , yields estimates of the proportion of bacterial, non-bacterial, and fossil CH_4 source strengths [Quay *et al.*, 1991]. We estimate that bacterial CH_4 sources contribute $\sim 70\%$, fossil CH_4 sources $\sim 20\%$, and biomass burning $\sim 10\%$ of the total CH_4 input.

Although the seasonal cycle in $\delta^{13}\text{C}$ dominates the time-series measurements in the northern hemisphere, there is evidence for a slight interannual increase up until 1992. Measurements at all four time-series locations indicate an increase in $\delta^{13}\text{C}$ and, when combined and area weighted, yield an average rate of approximately $0.04 \pm 0.02\text{‰}$ per year. A $\delta^{13}\text{C}$ increase indicates the ratio of bacterial to non-bacterial CH_4 source strength is decreasing. In 1993, however, there has been a leveling of the $\delta^{13}\text{C}$ trend at the northern sites and a decrease at SMO. This trend is similar to the recent $\delta^{13}\text{C}$ decrease observed at Baring Head, New Zealand (41°S) and Scott Base, Antarctica (78°) by Lowe *et al.* [1994], who attribute the decrease to a reduction of the CH_4 released from biomass burning. Measuring the interannual change in the $\delta^{13}\text{C}$ of atmospheric CH_4 at several sites over the longer term will be a useful indicator of changes in the proportion of




Fig. 1. The time series of the $\delta^{13}\text{C}$ of atmospheric CH_4 measured at BRW, Olympic Peninsula, MLO, and SMO since 1988.

bacterial and nonbacterial CH_4 inputs. These isotopic records should help us determine why the rate of methane increase has slowed so dramatically over the last few years.

REFERENCES

- Dlugokencky, E.J., K.A. Maserie, P.M. Lang, P.P. Tans, L.P. Steele and E.G. Nisbet, A dramatic decrease in the growth rate of atmospheric methane in the northern hemisphere during 1992, *Geophys. Res. Lett.*, **21**, 45-48, 1994.
- Cantrell, C.A., R.E. Shetter, A.H. McDaniel, J.G. Calvert, J.A. Davidson, D.C. Lowe, S.C. Tyler, R.J. Cicerone and J.P. Greenberg, Carbon kinetic isotope effect in the oxidation of methane by the hydroxyl radical, *J. Geophys. Res.*, **95**, 22,455-22,462, 1990.
- Lowe, D.C., C.A.M. Brenninkmeijer, G.W. Brailsford, K.R. Lassey, A.J. Gomez and E.G. Nisbet, Concentration and ^{13}C records of atmospheric methane in New Zealand and Antarctica: evidence for changes in methane sources, *J. Geophys. Res.*, **99**, 16,913-16,920, 1994.
- Quay, P.D., S.L. King, J. Stutsman, D.O. Wilbur, L.P. Steele, I. Fung, R.H. Gammon, T.A. Brown, G.W. Farwell, P.M. Groottes, and F.H. Schmidt, Carbon isotopic composition of atmospheric methane: Fossil and biomass burning source strengths, *Global Biogeochem. Cycles*, **5**, 25-47, 1991.
- Stevens, C.M. and F.E. Rust, The carbon isotopic composition of atmospheric methane, *J. Geophys. Res.*, **87**, 4879-4882, 1982.
- Wigley, T.M.L. and S.C.B. Raper, Implications for climate and sea level of revised IPCC emission scenarios, *Nature*, **357**, 293-300, 1992.